

CORRECTION

Open Access



Correction to: Disruption of hypoxia-inducible fatty acid binding protein 7 induces beige fat-like differentiation and thermogenesis in breast cancer cells

Masahiro Kawashima^{1,5*}, Karim Bensaad¹, Christos E. Zois¹, Alessandro Barberis², Esther Bridges¹, Simon Wigfield¹, Christoffer Lagerholm³, Ruslan I. Dmitriev^{4,6}, Mariko Tokiwa⁵, Masakazu Toi⁵, Dmitri B. Papkovsky⁴, Francesca M. Buffa² and Adrian L. Harris¹

Correction to: *Cancer Metab* 8, 13 (2020)

<https://doi.org/10.1186/s40170-020-00219-4>

Following publication of the original article [1], the authors identified an error in Fig. 7. The correct figure is given below.

thermogenesis in breast cancer cells. *Cancer Metab.* 2020;8:13 <https://doi.org/10.1186/s40170-020-00219-4>.

Author details

¹Department of Oncology, Molecular Oncology Laboratories, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford OX3 9DS, UK. ²Department of Oncology, Computational Biology and Integrative Genomics Lab, CRUK/MRC Institute for Radiation Oncology, University of Oxford, Old Road Campus Research Building, Roosevelt Drive, Oxford OX3 7DQ, UK. ³Wolfson Imaging Centre, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford OX3 9DS, UK. ⁴School of Biochemistry and Cell Biology, University College Cork, Cavanagh Pharmacy Building, 1.28, College Road, Cork, Ireland. ⁵Department of Breast Surgery, Graduate School of Medicine, Kyoto University, 54 Shogoin-Kawahara-cho, Sakyo-ku, Kyoto 606 8507, Japan. ⁶Institute for Regenerative Medicine, I.M. Sechenov First Moscow State University, Moscow, Russian Federation.

Published online: 10 August 2020

Reference

1. Kawashima M, Bensaad K, Zois CE, et al. Disruption of hypoxia-inducible fatty acid binding protein 7 induces beige fat-like differentiation and

The original article can be found online at <https://doi.org/10.1186/s40170-020-00219-4>.

* Correspondence: masa0123@kuhp.kyoto-u.ac.jp

¹Department of Oncology, Molecular Oncology Laboratories, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford OX3 9DS, UK

⁵Department of Breast Surgery, Graduate School of Medicine, Kyoto University, 54 Shogoin-Kawahara-cho, Sakyo-ku, Kyoto 606 8507, Japan

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 7 FABP7 knockdown (FABP7-Kd) increased cellular temperature. a Calibration curve of T probe generated using control cells (Ctrl). X and y axes show incubator temperatures and average fluorescence lifetime, respectively. b Average fluorescence lifetimes of Ctrl and FABP7-Kd. c Calculated cellular temperature of Ctrl (37°C) and FABP7-Kd. d Representative images of fluorescence lifetime imaging microscopy. Color scale indicates estimated temperature. Scale bars; 20 μm. Asterisks indicate nuclei locations. Error bars, SD; ***p* < 0.01, *n* = 3